arXiv:1205.2956v1 [math.GT] 14 May 2012

THE BOUNDARY OF A FIBERED FACE OF THE MAGIC 3-MANIFOLD AND THE ASYMPTOTIC BEHAVIOR OF THE MINIMAL PSEUDO-ANOSOVS DILATATIONS

EIKO KIN AND MITSUHIKO TAKASAWA

ABSTRACT. Let $\delta_{g,n}$ be the minimal dilatation of pseudo-Anosovs defined on an orientable surface of genus g with n punctures. Tsai proved that for any fixed $g \geq 2$, the logarithm of the minimal dilatation $\log \delta_{g,n}$ is on the order of $\frac{\log n}{n}$. The main result of this paper is that if $2g+1$ is relatively prime to s or $s+1$ for each $0 \leq s \leq g$, then

$$\limsup_{n \to \infty} \frac{n \log \delta_{g,n}}{\log n} \leq 2.$$

This improves the upper bound $\limsup\limits_{n\to\infty} \frac{n \log \delta_{g,n}}{\log n} \leq 2(2g+1)$ obtained from Tsai's examples. The examples of pseudo-Anosovs ϕ's which provide our bound above have the following property: The mapping torus M_ϕ of ϕ is a single hyperbolic 3-manifold N called the magic manifold, or the fibration of M_ϕ comes from a fibration of N by Dehn filling cusps along the boundary slopes of a fiber. The main tool in this paper is the boundary of a fibered face of N.

1. INTRODUCTION

Let $\Sigma = \Sigma_{g,n}$ be an orientable surface of genus g with n punctures and $\mathrm{Mod}(\Sigma)$ the mapping class group of Σ. According to the work of Nielsen and Thurston, elements of $\mathrm{Mod}(\Sigma)$ are classified into three types: periodic, reducible, pseudo-Anosov, see [2, 18]. The last one, pseudo-Anosov mapping classes have rich dynamical properties. The hyperbolization theorem by Thurston [21] relates the dynamics of pseudo-Anosovs and the geometry of hyperbolic fibered 3-manifolds. The theorem asserts that $\phi \in \mathrm{Mod}(\Sigma)$ is pseudo-Anosov if and only if the mapping torus M_ϕ of ϕ is a hyperbolic 3-manifold with finite volume.

Each pseudo-Anosov element $\phi \in \mathrm{Mod}(\Sigma)$ has a representative $\Phi : \Sigma \to \Sigma$ called a pseudo-Anosov homeomorphism. Such a homeomorphism is equipped with a constant $\lambda = \lambda(\Phi) > 1$ called the *dilatation* of Φ. This λ does not depend on the choice of a pseudo-Anosov homeomorphism $\Phi \in \phi$, and hence the *dilatation* $\lambda(\phi)$ of ϕ is defined to be $\lambda(\Phi)$. We call the quantities $\mathrm{ent}(\phi) = \log \lambda(\phi)$ and $\mathrm{Ent}(\phi) = |\chi(\Sigma)| \log \lambda(\phi)$ the *entropy* and *normalized entropy* of ϕ respectively, where $\chi(\Sigma)$ is the Euler characteristic of Σ.

If we fix Σ, the set of entropies of pseudo-Anosovs on Σ,

$$\{\mathrm{ent}(\phi) \mid \text{pseudo-Anosov } \phi \in \mathrm{Mod}(\Sigma)\},$$

is a closed discrete subset of $\mathbb{R}$, see [8]. In particular there exists a minimal dilatation, denoted by $\delta(\Sigma) > 1$, of pseudo-Anosov elements in $\mathrm{Mod}(\Sigma)$. One of the problems on the study of dynamics of pseudo-Anosovs is

---

Date: March 3, 2013.

2000 *Mathematics Subject Classification.* Primary 57M27, 37E30, Secondary 37B40 .

Key words and phrases. mapping class group, pseudo-Anosov, dilatation, entropy, magic manifold .

The first author is partially supported by Grant-in-Aid for Young Scientists (B) (No. 20740031), MEXT, Japan.

Problem 1.1 (Minimal dilatation problem)**.** *Determine the explicit value of* $\delta(\Sigma)$*. Describe pseudo-Anosov elements in* $\mathrm{Mod}(\Sigma)$ *which achieve* $\delta(\Sigma)$*.*

Problem 1.1 has wide interest in the following reason. The group $\mathrm{Mod}(\Sigma)$ acts properly discontinuously by isometries on the Teichmüller space $\mathrm{Teich}(\Sigma)$. The closed geodesics in the moduli space $\mathcal{M}(\Sigma) = \mathrm{Teich}(\Sigma)/\mathrm{Mod}(\Sigma)$ correspond precisely to the conjugacy classes of pseudo-Anosov elements of $\mathrm{Mod}(\Sigma)$. Furthermore the length of a closed geodesic associated to a pseudo-Anosov mapping class ϕ is equal to $\mathrm{ent}(\phi)(= \log \lambda(\phi))$. Hence Problem 1.1 is equivalent to determine the minimal closed geodesic in $\mathcal{M}(\Sigma)$ and to describe pseudo-Anosov elements in $\mathrm{Mod}(\Sigma)$ which achieve this minimum.

The minimal dilatations have been computed in several cases where $|\chi(\Sigma)|$ is small [3, 5, 13, 14]. However Problem 1.1 is widely open for most surfaces. Let us set $\delta_{g,n} = \delta(\Sigma_{g,n})$ and $\delta_g = \delta_{g,0}$. [1]Penner proved in [17] that $\log \delta_g \asymp \frac{1}{g}$. This work by Penner was a starting point on the study of the asymptotic behavior of the minimal dilatations on surfaces varying topology. Later it was proved by Hironaka-Kin [7] that $\log \delta_{0,n} \asymp \frac{1}{n}$, and by Tsai [22, 23] that $\log \delta_{1,n} \asymp \frac{1}{n}$. Recently Valdivia showed in [24] that given a rational number $r > 0$, $\log \delta_{g,n} \asymp \frac{1}{|\chi(\Sigma_{g,n})|}$, where (g, n) lies on the ray through the origin with the slope r. These results can be shown by Penner's lower bound

$$\frac{\log 2}{12g - 12 + 4n} \leq \log \delta_{g,n}$$

(see [17]) together with finding suitable families of pseudo-Anosovs.

The following remarkable theorem, due to Tsai, is in contrast with the cases of genus 0 or 1.

Theorem 1.2 ([22])**.** *For any fixed* $g \geq 2$*,* $\log \delta_{g,n} \asymp \frac{\log n}{n}$*.*

This paper concerns the following question which is motivated by Theorem 1.2.

Question 1.3. *Given* $g \geq 2$*, does* $\lim\limits_{n\to\infty} \dfrac{n \log \delta_{g,n}}{\log n}$ *exist? What is its value?*

This is an analogous question, posed by McMullen, which is asking whether $\lim\limits_{g\to\infty} g \log \delta_g$ exists or not, see [15].

Given $g \geq 2$, let $\{\psi_n \in \mathrm{Mod}(\Sigma_{g,n})\}_{n\in\mathbb{N}}$ be a family of pseudo-Anosovs. We say that the dilatation of the family $\{\psi_n\}$ *has the best possible asymptotic behavior* if $\log \lambda(\psi_n) \asymp \frac{\log n}{n}$. To best our knowledge, families of pseudo-Anosovs given by Tsai [22] are the only known examples whose dilatations have the best possible asymptotic behavior. In Appendix A, we review her constructions of pseudo-Anosovs, and show that her examples provide the upper bound $\limsup\limits_{n\to\infty} \frac{n \log \delta_{g,n}}{\log n} \leq 2(2g+1)$ (Proposition A.1). The purpose of this paper is to find another families of pseudo-Anosovs whose dilatations have the best possible asymptotic behavior, and to give a uniform upper bound of $\frac{n \log \delta_{g,n}}{\log n}$, independent of g, for infinitely many g's.

Our first result is

Theorem 1.4. *Given* $g \geq 2$*, there exists a sequence* $\{n_i\}_{i=0}^{\infty}$ *with* $n_i \to \infty$ *such that*

$$\limsup_{i\to\infty} \frac{n_i \log \delta_{g,n_i}}{\log n_i} \leq 2.$$

---

[1] Let A_g and B_g be functions on g. We write $A_g \asymp B_g$ if there exists a constant c, independent of g, such that $\frac{A_g}{c} < B_g < cA_g$.

To state the second result, we define the polynomial $B_{(g,p)}(t)$ for nonnegative integers g and p as follows.

$$B_{(g,p)}(t) = t^{2p+1}(t^{2g+1} - 1) - 2t^{p+g+1} - t^{2g+1} + 1.$$

We shall see that it has a unique real root, say $r_{(g,p)}$, greater than 1. The root $r_{(g,p)}$ satisfies

$$\lim_{p \to \infty} \frac{p \log r_{(g,p)}}{\log p} = 1$$

(see Lemma 4.3). Our second result is the following upper bound of the minimal dilatations.

Theorem 1.5. *For $g \geq 2$ and $p \geq 0$, suppose that* $\gcd(2g+1, p+g+1) = 1$. *Then*

$$\delta_{g,2p+i} \leq r_{(g,p)} \quad \textit{for each } i \in \{1,2,3,4\}.$$

See Table 1 for actual values of $r_{(g,p)}$'s.

Our last result is

Theorem 1.6. *Suppose that $g \geq 2$ satisfies*

$(*)$ $\gcd(2g+1, s) = 1$ *or* $\gcd(2g+1, s+1) = 1$ *for each* $0 \leq s \leq g$.

Then

$$\limsup_{n \to \infty} \frac{n \log \delta_{g,n}}{\log n} \leq 2.$$

For example $(*)$ holds for $g = 4$ since 9 is relatively prime to $1, 2, 4$ and 5, but $(*)$ does not hold for $g = 7$ because $\gcd(15, 5) = 5$ and $\gcd(15, 6) = 3$. We would like to point out that infinitely many g's satisfy $(*)$. In fact if $2g+1$ is prime, then $2g+1$ is relatively prime to s' for each $1 \leq s' \leq g+1$. Such a g enjoys $(*)$, and this leads to

Corollary 1.7. *If $2g+1$ is prime for $g \geq 2$, then*

$$\limsup_{n \to \infty} \frac{n \log \delta_{g,n}}{\log n} \leq 2.$$

Remark 1.8. *One can simplify $(*)$ since $2g+1$ is always relative prime to $1, 2$ and g. In the case $g \geq 5$, $(*)$ is equivalent to*

$(**)$ $\gcd(2g+1, s) = 1$ *or* $\gcd(2g+1, s+1) = 1$ *for each* $3 \leq s \leq g-2$.

Our results are proved by using the theory on fibered faces of hyperbolic fibered 3-manifolds M, developed by Thurston and Fried, see Section 3. Let $\| \cdot \| : H_2(M, \partial M; \mathbb{R}) \to \mathbb{R}$ be the Thurston norm, and let Ω be a fibered face of M. (Fibered faces of M are some of the top dimensional faces of the unit ball U_M with respect to the norm $\| \cdot \|$.) The work of Thurston in [19] tells us that if M has the second Betti number more than 1, then it admits a family of fibrations on M dominated by $int(C_\Omega)$, where C_Ω is the core over Ω with the origin and $int(C_\Omega)$ is its interior. In other words, such a fibered 3-manifold provides infinitely many pseudo-Anosovs defined on surfaces with variable topology. By the work of Fried [4], the entropy function defined on these fibrations admits a unique continuous extension $\mathrm{ent} : int(C_\Omega) \to \mathbb{R}$. The entropy function, ent, has the property that $\frac{1}{\mathrm{ent}} : int(C_\Omega) \to \mathbb{R}$ is concave. Moreover the restriction $\mathrm{ent}|_{int(\Omega)} : int(\Omega) \to \mathbb{R}$ enjoys that $\mathrm{ent}(a)$ goes to ∞ as $a \in int(\Omega)$ goes to a point on the boundary of Ω, see [4].

These properties give us the following useful observation: For any compact set $\mathcal{D} \subset int(\Omega)$, there exists a constant $C = C_{\mathcal{D}} > 0$ satisfying the following. Let $a \in int(C_\Omega)$ be any integral class of $H_2(M, \partial M; \mathbb{Z})$ and let Φ_a be the monodromy of the fibration associated to a. Then the normalized entropy $\mathrm{Ent}(\Phi_a) \leq C$ whenever $\overline{a} \in \mathcal{D}$, where $\overline{a}$ is the projective class of a.

This observation enables us to investigate the asymptotic behavior of the minimal dilatations on surfaces varying topology. The following theorem, showing the best known upper bounds, are proved by using a technique with the same flavor as the observation above.

Theorem 1.9 ([7, 11] for (1), [9] for (2), [6, 1, 12] for (3)). *Let D_n be an n-punctured disk.*

1. $\limsup_{n\to\infty} n \log \delta_{0,n} \leq 2\log(2+\sqrt{3})$.
2. $\limsup_{n\to\infty} n \log \delta_{1,n} \leq 2\log \delta(D_4) \approx 1.6628$, *where* $\delta(D_4) \approx 2.2966$ *is the largest real root of* $t^4 - 2t^3 - 2t + 1$.
3. $\limsup_{g\to\infty} g \log \delta_g \leq \log \delta(D_3) = \log \delta(\Sigma_{1,1}) = \log(\frac{3+\sqrt{5}}{2})$.

However for any fixed $g \geq 2$, the same technique doesn't work in order to give an upper bound of $\delta_{g,n}$ varying n because of Theorem 1.2. Theorem 1.2 implies that there exists no constant $C > 0$, independent of n, such that the minimal normalized entropy $|\chi(\Sigma_{g,n})| \log \delta_{g,n}$ satisfies $|\chi(\Sigma_{g,n})| \log \delta_{g,n} < C$. Thus if there exists a sequence of integral classes $\{a_i\}$ with $a_i \in int(C_\Omega)$ such that the fiber of the fibration associated to a_i is a surface of genus g and n_i boundary components with $n_i \to \infty$, then the accumulation points of the sequence of projective classes $\{\overline{a_i}\}$ must lie on the boundary of Ω. (This is because there exists no constant $C > 0$, independent of i, such that $\mathrm{Ent}(\Phi_{a_i}) \leq C$, see Theorem 1.2.)

Nevertheless we focus on a particular hyperbolic fibered 3-manifold, called the *magic manifold* N. This is the exterior of the 3 chain link $\mathcal{C}_3$, see Figure 1. It turns out that the boundary of a fibered face of N is a useful tool for Question 1.3.

Our examples of pseudo-Anosovs ϕ's which provide the bounds in Theorems 1.4 and 1.6 have the following property: The mapping torus M_ϕ of ϕ is homeomorphic to N, or the fibration of M_ϕ comes from a fibration of N by Dehn filling cusps along the boundary slopes of a fiber.



FIGURE 1. (left) 3 chain link $\mathcal{C}_3$ (right) link whose exterior is homeomorphic to N. (This is an illustration of a fibration on N with the fiber homeomorphic to the 4-punctured sphere.)

Now we turn our attention to the minimal volume problems on hyperbolic 3-manifolds. The set of volumes of hyperbolic 3-manifolds is a well-ordered closed subset in $\mathbb{R}$ of order type ω^ω, see [18]. In particular if we fix a surface Σ, then there exists a minimum among volumes of hyperbolic Σ-bundles over the circle. It is known that hyperbolic volume decreases after Dehn filling, see [16, 18]. By appealing to this result on the surface bundles over the circle in the proofs of Theorems 1.4, 1.6, we obtain

Proposition 1.10. *Given $g \geq 2$, there exists a sequence $\{n_i\}_{i=0}^{\infty}$ with $n_i \to \infty$ such that the minimal volume of Σ_{g,n_i}-bundles over the circle is less than or equal to the volume of the magic manifold $N \approx 5.3334$. Furthermore if $g \geq 2$ satisfies $(*)$, then for any $n \geq 3$, the minimal volume of $\Sigma_{g,n}$-bundles over the circle is less than or equal to the volume of N.*

We close the introduction by asking

Question 1.11 (cf. Theorems 1.4 and 1.6)**.** *Does there exists a universal constant* $C > 0$ *such that given* $g \geq 2$, $\limsup_{n\to\infty} \frac{n \log \delta_{g,n}}{\log n} \leq C$ *holds? If this is true, then does* $\limsup_{n\to\infty} \frac{n\log\delta_{g,n}}{\log n} \leq 2$ *hold?*

2. Roots of polynomials

This section concerns the asymptotic behavior of roots of families of polynomials. We will use the claims given in this section to analyze the asymptotic behavior of the dilatations in families of fibered classes.

Let $g(t) = a_n t^{b_n} + a_{n-1}t^{b_{n-1}} + \cdots + a_0 t^{b_0}$ be a polynomial with nonzero real coefficients a_i, where b_i's are integers satisfying $b_n > b_{n-1} > \cdots > b_1 > b_0 \geq 0$. Let $\mathfrak{D}(g)$ be the number of variations in sign in the sequence $a_n, a_{n-1}, \cdots, a_0$ of the coefficients. For example if $g(t) = t^4 + t^3 - 2t^2 + t - 1$, then $\mathfrak{D}(g) = 3$.

Theorem 2.1 ([25])**.** *Let* $g(t)$ *be a polynomial as above. Then the number of positive real roots of* $g(t)$ *(counted with multiplicities) is either equal to* $\mathfrak{D}(g)$ *or less than* $\mathfrak{D}(g)$ *by an even integer.*

We fix integers $r \geq 0$, $s > 0$, $u > 0$, and fix a polynomial $Q(t)$ whose coefficients are positive integers. Consider the following polynomial $P_m(t)$ for each $m \in \mathbb{N}$:

$$P_m(t) = t^{2m+r}(t^s - 1) + 1 - Q(t)t^m - t^u = t^{2m+r+s} - t^{2m+r} - Q(t)t^m - t^u + 1.$$

Proposition 2.2.

(1) *Suppose that the degree of each monomial other than* t^{2m+r+s} *is less than that of* t^{2m+r+s}*, that is*

$$2m + r + s > \max\{\deg(Q(t)t^m), u\},$$

where $\deg(f)$ *is the degree of the polynomial* f*. Then* $P_m(t)$ *has a unique real root* λ_m *greater than* 1*.*

(2) *Given* $0 < c_1 < 1$ *and* $c_2 > 1$*, there exists* $L > 0$ *such that*

$$m^{\frac{c_1}{m}} < \lambda_m < m^{\frac{c_2}{m}} \quad \textit{for any } m > L.$$

(3) *We have*

$$\lim_{m\to\infty} \frac{m \log \lambda_m}{\log m} = 1.$$

Proof. (1) Under the assumption on the degree in (1), we have $\mathfrak{D}(P_m) = 2$. By Theorem 2.1, the number of positive real roots of P_m is exactly equal to 2 since $P_m(0) = 1$, $P_m(1) < 0$. Because $P_m(t)$ goes to ∞ as t does, $P_m(t)$ has a unique real root $\lambda_m > 1$.

(2) We have

$$P_m(t)t^{-(2m+r)} = (t^s - 1) + t^{-(2m+r)} - Q(t)t^{-(m+r)} - t^{-(2m+r-u)}.$$

Put $f_m(t)$ and $g_m(t)$ as follows.

$$\begin{aligned} f_m(t) &= (t^s - 1) + t^{-(2m+r)}, \\ g_m(t) &= Q(t)t^{-(m+r)} + t^{-(2m+r-u)}. \end{aligned}$$

It is enough to prove that if we take $0 < c_1 < 1$ and $c_2 > 1$, then $f_m(m^{\frac{c_1}{m}}) < g_m(m^{\frac{c_1}{m}})$ and $f_m(m^{\frac{c_2}{m}}) > g_m(m^{\frac{c_2}{m}})$ for m large.

If $t = m^{\frac{c}{m}}$ for some $c > 0$, then

$$t^s = (m^{\frac{c}{m}})^s = \big((e^{\log m})^{\frac{c}{m}}\big)^s = (e^{\frac{c\log m}{m}})^s = e^{\frac{sc\log m}{m}}.$$

By Maclaurin expansion of the exponential function, we have

$$e^{\frac{sc\log m}{m}} = 1 + \frac{sc\log m}{m} + R_2,$$

where $R_2 = \frac{e^w}{2}(\frac{sc\log m}{m})^2$ for some $0 < w < \frac{sc\log m}{m}$. Since $\frac{sc\log m}{m}$ goes to 0 as m goes to ∞, we may assume that $\frac{e^w}{2} < B$ for some constant $B > 0$. Then

$$\begin{array}{rcl} f_m(m^{\frac{c}{m}}) & = & (m^{\frac{c}{m}})^s - 1 + (m^{\frac{c}{m}})^{-(2m+r)} \\ & = & \dfrac{sc\log m}{m} + R_2 + \dfrac{m^{1-c(2+\frac{r}{m})}}{m} \\ & < & \dfrac{sc\log m}{m} + B(\dfrac{sc\log m}{m})^2 + \dfrac{m^{1-c(2+\frac{r}{m})}}{m} \\ & = & \dfrac{sc\log m}{m} + Bs^2c^2(\dfrac{\log m}{m})^2 + \dfrac{m^{1-c(2+\frac{r}{m})}}{m} \\ & < & \dfrac{sc\log m}{m} + Bs^2c^2(\dfrac{\log m}{m}) \cdot 1 + \dfrac{m^{1-c(2+\frac{r}{m})}}{m} \\ & = & \dfrac{(sc + Bs^2c^2)\log m + m^{1-c(2+\frac{r}{m})}}{m}. \end{array}$$

Note that from the second equality above, we have

$$f_m(m^{\frac{c}{m}}) > \frac{sc\log m}{m}. \tag{2.1}$$

If we write $Q(t) = \sum_{j=0}^{\ell} a_j t^j$, where a_j is a nonnegative integer, then

$$\begin{array}{rcl} g_m(m^{\frac{c}{m}}) & = & Q(m^{\frac{c}{m}})m^{-c(1+\frac{r}{m})} + m^{-c(2+\frac{r}{m}-\frac{u}{m})} \\ & = & Q(m^{\frac{c}{m}})\dfrac{m^{1-c(1+\frac{r}{m})}}{m} + \dfrac{m^{1-c(2+\frac{r}{m}-\frac{u}{m})}}{m} \\ & = & \displaystyle\sum_{j=0}^{\ell}\Big(\dfrac{a_j m^{1-c(1+\frac{r}{m}-\frac{j}{m})}}{m}\Big) + \dfrac{m^{1-c(2+\frac{r}{m}-\frac{u}{m})}}{m}. \\ & = & \dfrac{\big(\sum_{j=0}^{\ell} a_j m^{1-c(1+\frac{r}{m}-\frac{j}{m})}\big) + m^{1-c(2+\frac{r}{m}-\frac{u}{m})}}{m}. \end{array}$$

Suppose that the above $c > 0$ satisfies $0 < c < \frac{1}{2}$. The four terms $\log m$, $m^{1-c(2+\frac{r}{m})}$, $m^{1-c(1+\frac{r}{m}-\frac{j}{m})}$ and $m^{1-c(2+\frac{r}{m}-\frac{u}{m})}$ go to ∞ as m does, since the last three terms have the positive power of m. Notice that given $C > 0$, there exists a constant $L_C > 0$ such that $\log m < m^C$ for any $m > L_C$. Observe that among the four terms above, $m^{1-c(1+\frac{r}{m}-\frac{j}{m})}$ is dominant because

$$1 - c(1 + \frac{r}{m} - \frac{j}{m}) > 1 - c(2 + \frac{r}{m} - \frac{u}{m}) \geq 1 - c(2 + \frac{r}{m}).$$

These imply that $f_m(m^{\frac{c_1}{m}}) < g_m(m^{\frac{c_1}{m}})$ holds for m large.

Suppose that $\frac{1}{2} \leq c < 1$. We can check that $m^{1-c(1+\frac{r}{m}-\frac{j}{m})}$ is still dominant among four terms. (Some of the terms except $m^{1-c(1+\frac{r}{m}-\frac{j}{m})}$ may be bounded as m goes to ∞.) Therefore $f_m(m^{\frac{c_1}{m}}) < g_m(m^{\frac{c_1}{m}})$ holds for m large.

Finally consider the case $c > 1$. Then $m^{1-c(2+\frac{r}{m})}$, $m^{1-c(1+\frac{r}{m}-\frac{j}{m})}$ and $m^{1-c(2+\frac{r}{m}-\frac{u}{m})}$ go to 0 as m goes to ∞, because these terms have the negative power of m as m goes to ∞. Thus the numerator of $g_m(m^{\frac{c}{m}})$ goes to 0 as m tends to ∞. On the other hand, $f_m(m^{\frac{c}{m}}) > \frac{sc\log m}{m}$ holds (see (2.1)), and hence the numerator of $f_m(m^{\frac{c}{m}}) = \frac{sc\log m + mR_2 + m^{1-c(2+\frac{r}{m})}}{m}$ goes to ∞ as m does. Thus $f_m(m^{\frac{c_1}{m}}) > g_m(m^{\frac{c_1}{m}})$ for m large.

(3) By the inequality in (2), we have $c_1 < \frac{m\log\lambda_m}{\log m} < c_2$ for m large. Since $0 < c_1 < 1$ and $c_2 > 1$ are any numbers, $\frac{m\log\lambda_m}{\log m}$ goes to 1 as m goes to ∞. □

Lemma 2.3. *Let* $\{\kappa_m\}_{m\in\mathbb{N}}$ *be a sequence of real numbers such that* $\kappa_m > 1$ *for each* m. *Suppose that given* $0 < c_1 < 1$ *and* $c_2 > 1$, *there exists* $L > 0$ *such that*

$$m^{\frac{c_1}{m}} < \kappa_m < m^{\frac{c_2}{m}} \quad \textit{for any } m > L.$$

Then for any fixed integers $q > 0$ *and* $r \geq 0$, *we have*

$$\lim_{m\to\infty} \frac{(qm+r)\log\kappa_m}{\log(qm+r)} = q.$$

Proof. By the assumption,

$$\frac{c_1\log m}{m} < \log\kappa_m < \frac{c_2\log m}{m}.$$

Let us set $n = qm + r$. We substitute $m = \frac{n-r}{q}$ for the inequality above:

$$\frac{c_1\log\left(\frac{n-r}{q}\right)}{\frac{n-r}{q}} < \log\kappa_m < \frac{c_2\log\left(\frac{n-r}{q}\right)}{\frac{n-r}{q}}.$$

Hence

$$\frac{qc_1\big(\log(n-r)-\log q\big)}{n-r} < \log\kappa_m < \frac{qc_2\big(\log(n-r)-\log q\big)}{n-r}.$$

If each side is multiplied by $\frac{n}{\log n}$, then we obtain

$$\frac{qc_1\big(\log(n-r)-\log q\big)n}{(n-r)\log n} < \frac{n\log\kappa_m}{\log n} < \frac{qc_2\big(\log(n-r)-\log q\big)n}{(n-r)\log n}.$$

Note that $\frac{\big(\log(n-r)-\log q\big)n}{(n-r)\log n} = \big(\frac{n}{n-r}\big)\big(\frac{\log(n-r)-\log q}{\log n}\big)$ goes to 1 as n goes to ∞. Since $0 < c_1 < 1$ and $c_2 > 1$ be any numbers, it follows that

$$\lim_{m\to\infty} \frac{(qm+r)\log\kappa_m}{\log(qm+r)} = q.$$

□

3. Thurston norm and fibered 3-manifolds

Let M be an oriented hyperbolic 3-manifold with boundary ∂M (possibly $\partial M = \emptyset$). First we recall the Thurston norm $\| \cdot \| : H_2(M, \partial M; \mathbb{R}) \to \mathbb{R}$, and recall the relation between $\| \cdot \|$ and fibrations on M when M fibers over the circle. (See [19] for more details.)

The Thurston norm $\| \cdot \|$ is defined for an integral class $a \in H_2(M, \partial M; \mathbb{Z})$ by

$$\|a\| = \min_F \{-\chi(F)\},$$

where the minimum is taken over all oriented surface F embedded in M, satisfying $a = [F]$, with no components of non-negative Euler characteristic. The surface F which realizes this minimum is called the *minimal representative* of a, denoted by F_a. The norm $\| \cdot \|$ defined on integral classes admits a unique continuous extension $\| \cdot \| : H_2(M, \partial M; \mathbb{R}) \to \mathbb{R}$ which is linear on the ray through the origin. The unit ball U_M with respect to the Thurston norm is a compact, convex polyhedron.

Suppose that M is a surface bundle over the circle and let F be its fiber. The fibration determines a cohomology class $a^* \in H^1(M; \mathbb{Z})$, and hence a homology class $a \in H_2(M, \partial M; \mathbb{Z})$ by Poincaré duality. Thurston proved in [19] that there exists a top dimensional face Ω on ∂U_M such that $a = [F]$ is an integral class of $int(C_\Omega)$. On the other hand, the minimal representative F_a for any integral class a in $int(C_\Omega)$ becomes a fiber of the fibration associated to a. Such a face Ω is called a *fibered face*, and an integral class $a \in int(C_\Omega)$ is called a *fibered class*.

The set of integral and rational classes of $int(C_\Omega)$ are denoted by $int(C_\Omega(\mathbb{Z}))$ and $int(C_\Omega(\mathbb{Q}))$ respectively. When $a \in int(C_\Omega(\mathbb{Z}))$ is primitive, the associated fibration on M has a connected fiber represented by F_a. Let $\Phi_a : F_a \to F_a$ be the monodromy. Since M is hyperbolic, $\phi_a = [\Phi_a]$ is pseudo-Anosov. The *dilatation* $\lambda(a)$ and *entropy* $\mathrm{ent}(a) = \log \lambda(a)$ are defined as the dilatation and entropy of ϕ_a respectively.

We now recall the work of Fried. The entropy defined on primitive fibered classes is extended to rational classes as follows: For a rational number r and a primitive fibered class a, the entropy $\mathrm{ent}(ra)$ is defined by $\frac{1}{|r|}\mathrm{ent}(a)$. Fried proved that $\frac{1}{\mathrm{ent}} : int(C_\Omega(\mathbb{Q})) \to \mathbb{R}$ is concave [4], and in particular $\mathrm{ent} : int(C_\Omega(\mathbb{Q})) \to \mathbb{R}$ admits a unique continuous extension

$$\mathrm{ent} : int(C_\Omega) \to \mathbb{R}.$$

Moreover, the restriction of ent to the open fibered face $int(\Omega)$ has the property that $\mathrm{ent}(a)$ goes to ∞ as $a \in int(\Omega)$ goes to a point on $\partial\Omega$. Thus we have a continuous function

$$\mathrm{Ent} = \| \cdot \| \, \mathrm{ent}(\cdot) : int(C_\Omega) \to \mathbb{R}$$

which is constant on each ray in $int(C_\Omega)$ through the origin.

4. The magic 3-manifold

The fibers/monodromies of fibrations on N have been studied in [11, 12]. In Sections 4.1 and 4.2 we recall some results from the papers. Section 4.2 includes the proofs of Theorems 1.4, 1.5 and 1.6.

Recall that $\Sigma_{g,n}$ is an orientable surface of genus g with n punctures. By abusing the notation, we sometimes denote an orientable surface of genus g with n boundary components by $\Sigma_{g,n}$ in the rest of the paper.

4.1. Fibered face.

Let K_α, K_β and K_γ be the components of the 3 chain link $\mathcal{C}_3$. They bound the oriented disks F_α, F_β and F_γ with 2 holes, see Figure 2(right). Let $\alpha = [F_\alpha]$, $\beta = [F_\beta]$, $\gamma = [F_\gamma] \in H_2(N, \partial N; \mathbb{Z})$. The set $\{\alpha, \beta, \gamma\}$ is a basis of $H_2(N, \partial N; \mathbb{Z})$. Figure 2(left) illustrates the

Thurston norm ball U_N for N which is the parallelepiped with vertices $\pm\alpha$, $\pm\beta$, $\pm\gamma$, $\pm(\alpha+\beta+\gamma)$, see [19].

Because of the symmetry of $\mathcal{C}_3$, every top dimensional face of U_N is a fibered face, and it is enough to consider a particular fibered face in order to study the monodromies of fibrations on N. We denote a class $x\alpha + y\beta + z\gamma \in H_2(N, \partial N)$ by (x, y, z). We pick a fibered face Δ with vertices $\alpha = (1, 0, 0)$, $\alpha+\beta+\gamma = (1, 1, 1)$, $\beta = (0, 1, 0)$ and $-\gamma = (0, 0, -1)$. The open face $int(\Delta)$ is written by

$$int(\Delta) = \{(X, Y, Z) \mid X + Y - Z = 1,\ X > 0,\ Y > 0,\ X > Z,\ Y > Z\}.$$

A class $a = (x, y, z) \in H_2(N, \partial N)$ is an element of $int(C_\Delta)$ if and only if $x > 0$, $y > 0$, $x > z$ and $y > z$. In this case $\|a\|$ is equal to $x + y - z$. Its projective class $\overline{a} = \overline{(x, y, z)} \in int(\Delta)$ is given by $\overline{a} = (\frac{x}{x+y-z}, \frac{y}{x+y-z}, \frac{z}{x+y-z})$. One of the coordinates, say $\frac{z}{x+y-z}$, is not needed to represent $\overline{a}$, so we denote $(\frac{x}{x+y-z}, \frac{y}{x+y-z}, \frac{z}{x+y-z})$ by $[\frac{x}{x+y-z}, \frac{y}{x+y-z}]$ for simplicity. With this notation

$$int(\Delta) = \{[X, Y] \mid 0 < X < 1,\ 0 < Y < 1\}.$$



FIGURE 2. (left) Thurston norm ball U_N. (right) F_α, F_β, F_γ. [arrows indicate the normal direction of oriented surfaces.]

Let $a = (x, y, z)$ be a primitive fibered class in $int(C_\Delta)$. The minimal representative of this class is denoted by F_a or $F_{(x,y,z)}$. We recall the formula which tells us that the number of the boundary components of F_a. We denote the tori $\partial\mathcal{N}(K_\alpha)$, $\partial\mathcal{N}(K_\beta)$, $\partial\mathcal{N}(K_\gamma)$ by T_α, T_β, T_γ respectively, where $\mathcal{N}(K)$ be a regular neighborhood of a knot K in S^3. Let us set $\partial_\alpha F_{(x,y,z)} = \partial F_{(x,y,z)} \cap T_\alpha$ which consists of the parallel simple closed curves on T_α. We define the subsets $\partial_\beta F_{(x,y,z)}$, $\partial_\gamma F_{(x,y,z)} \subset \partial F_{(x,y,z)}$ in the same manner. By [11, Lemma 3.1] the number of the boundary components $\sharp(\partial F_{(x,y,z)})$ is equal to

$$\gcd(x, y+z) + \gcd(y, z+x) + \gcd(z, x+y),$$

where $\gcd(0, w)$ is defined by $|w|$. More precisely

$$\sharp(\partial_\alpha F_{(x,y,z)}) = \gcd(x, y+z),\ \sharp(\partial_\beta F_{(x,y,z)}) = \gcd(y, z+x),\ \sharp(\partial_\gamma F_{(x,y,z)}) = \gcd(z, x+y).$$

Moreover the boundary slope of each component of $\partial_\alpha F_{(x,y,z)}$ equals $\frac{-(y+z)}{x}$. Similarly the boundary slope of each component of $\partial_\beta F_{(x,y,z)}$ (resp. $\partial_\gamma F_{(x,y,z)}$) is given by $\frac{-(z+x)}{y}$ (resp. $\frac{-(x+y)}{z}$).

We define slope(x, y, z) to be slope$(x, y, z) = (\frac{-(y+z)}{x}, \frac{-(z+x)}{y}, \frac{-(x+y)}{z})$.

4.2. **Dilatations of fibered classes.** Teichmüller polynomial P_Δ on the fibered face Δ tells us that the dilatation $\lambda_{(x,y,z)}$ of a fibered class $(x,y,z) \in int(C_\Delta)$ is the largest real root of

$$f_{(x,y,z)}(t) = t^{x+y-z} - t^x - t^y - t^{x-z} - t^{y-z} + 1,$$

see [11, Theorem 3.1]. Notice that $(x,y,z) \in int(C_\Delta)$ implies that $\mathfrak{D}(f_{(x,y,z)}) = 2$. By Theorem 2.1, $\lambda_{(x,y,z)}$ is indeed a unique real root greater than 1 of $f_{(x,y,z)}(t)$.

Let $\Phi_{(x,y,z)} : F_{(x,y,z)} \to F_{(x,y,z)}$ be the monodromy of the fibration associated to a primitive class $(x,y,z) \in int(C_\Delta)$ with a fiber $F_{(x,y,z)}$. Let $\mathcal{F}_{(x,y,z)}$ be the stable foliation of $\Phi_{(x,y,z)}$. The components of $\partial_\alpha F_{(x,y,z)}$ (resp. $\partial_\beta F_{(x,y,z)}$, $\partial_\gamma F_{(x,y,z)}$) are permuted cyclically by $\Phi_{(x,y,z)}$. In particular the number of prongs of $\mathcal{F}_{(x,y,z)}$ at a component of $\partial_\alpha F_{(x,y,z)}$ (resp. $\partial_\beta F_{(x,y,z)}$, $\partial_\gamma F_{(x,y,z)}$) is independent of the choice of the component. By [12], $\mathcal{F}_{(x,y,z)}$ is $\frac{x}{\gcd(x,y+z)}$-pronged at each component of $\partial_\alpha F_{(x,y,z)}$, $\frac{y}{\gcd(y,x+z)}$-pronged at each component of $\partial_\beta F_{(x,y,z)}$, $\frac{x+y-2z}{\gcd(z,x+y)}$-pronged at each component of $\partial_\gamma F_{(x,y,z)}$, and $\mathcal{F}_{(x,y,z)}$ has no singularities in the interior of $F_{(x,y,z)}$.

4.3. **Proofs of Theorems 1.4, 1.5 and 1.6.** We begin by defining linear sections in $H_2(N, \partial N; \mathbb{R})$. For $r \in \mathbb{Q}$ or $r = \infty$, define

$$S_\beta(r) = \{(x,y,z) \in H_2(N, \partial N; \mathbb{R}) \mid r = \tfrac{z+x}{-y}\}.$$

In the rest of this section, we restrict our attention to the section $S_\beta(-1)$. Figure 3 is an illustration of $S_\beta(-1) \cap U_N$ and $S_\beta(-1) \cap \Delta$. Since there is a symmetry of $S_\beta(-1) \cap U_N$, it is enough to study the fibered classes of $S_\beta(-1) \cap int(C_\Delta)$.



FIGURE 3. Linear sections. (left) $S_\beta(-1) \cap U_N$. (right) $S_\beta(-1) \cap \Delta$.

If we take $\mathfrak{a} = (1,1,0)$ and $\mathfrak{b} = (0,1,1)$, then $\{\mathfrak{a}, \mathfrak{b}\}$ becomes a basis of $S_\beta(-1) \cap H_2(N, \partial N; \mathbb{Z})$. One can check that $k\mathfrak{a} + \ell\mathfrak{b} = (k, k+\ell, \ell) \in int(C_\Delta)$ if and only if $k > 0$ and $-k < \ell < k$. Therefore

$$S_\beta(-1) \cap int(\Delta) = \{[\tfrac{1}{2}, Y] \mid 0 < Y < 1\}.$$

Lemma 4.1. *Suppose that $k\mathfrak{a} + \ell\mathfrak{b}$ is a primitive integral class of $int(C_\Delta)$.*

1. *$F_{k\mathfrak{a}+\ell\mathfrak{b}}$ is a surface of genus $\frac{k-\ell}{2}$ with $k+\ell+2$ boundary components if both k and ℓ are odd.*
2. *$F_{k\mathfrak{a}+\ell\mathfrak{b}}$ is a surface of genus $\frac{k-\ell-1}{2}$ with $k+\ell+3$ boundary components otherwise.*

Proof. The number of the boundary components of $F_{k\mathfrak{a}+\ell\mathfrak{b}}$ is equal to

$$\gcd(k, 2\ell) + k + \ell + \gcd(\ell, 2k).$$

It is equal to $k+\ell+2$ if both k and ℓ are odd. Otherwise $k+\ell+3$ (since k and ℓ are relatively prime). The genus of $F_{k\mathfrak{a}+\ell\mathfrak{b}}$ is computed from $|\chi(F_{k\mathfrak{a}+\ell\mathfrak{b}})| = \|k\mathfrak{a}+\ell\mathfrak{b}\| = 2k$ and the number of the boundary components. □

For $g \geq 0$ and $p \geq 0$, define a fibered class $a_{(g,p)}$ as follows.

$$a_{(g,p)} = (p+g+1)\mathfrak{a} + (p-q)\mathfrak{b} = (p+g+1, 2p+1, p-g) \in int(C_\Delta).$$

Note that $\|a_{(g,p)}\| = 2p+2g+2$. The class $a_{(g,p)}$ is primitive if and only if $2g+1$ and $p+g+1$ are relatively prime. One can check that

$$B_{(g,p)}(t) = f_{(p+g+1,2p+1,p-g)}(t)$$

(see Section 1 for the definition of $B_{(g,p)}(t)$), and hence the unique real root $r_{(g,p)} > 1$ of $B_{(g,p)}(t)$ equals the dilatation $\lambda(a_{(g,p)})$.

TABLE 1. $r_{(g,p)} = \lambda(a_{(g,p)})$. (Numbers with the round bracket in the table mean that in these cases, the fibered classes $a_{(g,p)}$'s are not primitive, i.e, $2g+1$ and $p+g+1$ are not relatively prime. If $a_{(g,p)}$ is primitive, then $\delta_{g,2p+i} \leq r_{(g,p)}$ for each $i \in \{1,2,3,4\}$, see Theorem 1.5.)

p	$r_{(2,p)}$	$r_{(3,p)}$	$r_{(4,p)}$	$r_{(5,p)}$	$r_{(6,p)}$	$r_{(7,p)}$
0	1.7220	1.5560	1.4579	1.3924	1.3450	1.3090
1	1.4012	1.3247	(1.2767)	1.2431	1.2181	(1.1986)
2	(1.3013)	1.2483	1.2144	1.1903	1.172	(1.1578)
3	1.2483	(1.2069)	1.1801	1.1608	1.1462	1.1346
4	1.2144	1.1801	(1.1575)	1.1413	1.1289	(1.1190)
5	1.1903	1.1608	1.1413	(1.1271)	1.1163	1.1076
6	1.1721	1.1462	1.1289	1.1163	(1.1066)	1.0988
7	(1.1578)	1.1346	(1.1190)	1.1076	1.0988	(1.0917)
8	1.1461	1.1251	1.1109	1.1004	1.0924	1.0859
9	1.1364	1.1171	1.1040	1.0944	1.0870	1.0809
10	1.1281	(1.1103)	(1.0982)	1.0893	1.0823	(1.0767)
11	1.1210	1.1045	1.0932	1.0848	1.0783	1.0730
12	(1.1148)	1.0993	1.0887	1.0809	1.0747	(1.0698)
13	1.1093	1.0948	(1.0848)	1.0774	1.0716	(1.0669)
14	1.1044	1.0907	1.0813	1.07428	1.0687	1.0643
15	1.1000	1.0870	1.0781	1.0714	1.0662	1.0619
16	1.0960	1.0837	(1.0752)	(1.0688)	1.0638	(1.0598)
17	(1.0924)	(1.0807)	1.0726	1.0665	1.0617	(1.0578)
18	1.0891	1.0779	1.0702	1.0643	1.0597	1.0560
19	1.0861	1.0754	(1.0679)	1.0624	(1.0579)	(1.0543)
20	1.0833	1.0730	1.0659	1.0605	1.0563	1.0528

The rational class $[\frac{1}{2}, \frac{2p+1}{2p+2g+2}] \in int(\Delta)$ represents the projective class $\overline{a_{(g,p)}}$ of $a_{(g,p)}$. This leads to the following. (See also Figure 4.)

Lemma 4.2. *If we fix $g \geq 0$, the sequence $\{\overline{a_{(g,p)}}\}_{p \in \mathbb{N}}$ converges to $[\frac{1}{2}, 1] \in \partial\Delta$ as p goes to ∞.*

By Proposition 2.2(2), we have the following.

Lemma 4.3. *We fix $g \geq 0$. Given $0 < c_1 < 1$ and $c_2 > 1$, there exists $L > 0$ such that*

$$p^{\frac{c_1}{p}} < r_{(g,p)} < p^{\frac{c_2}{p}} \quad \textit{for any } p > L.$$


Y
$\overline{a_{(0,p+1)}}$
enlarge
1
$\overline{a_{(0,p)}}$
3/4
$\overline{a_{(0,1)}}$
1/2
$\overline{a_{(0,0)}}$
0
1/2
1
X
$\overline{a_{(0,p+1)}} = \overline{a_{(g,(2g+1)(p+1)+g)}}$
$\overline{a_{(g,(2g+1)p+g+i)}}$
$\overline{a_{(g,(2g+1)p+g+1)}}$
$\overline{a_{(0,p)}} = \overline{a_{(g,(2g+1)p+g)}}$
X=1/2

FIGURE 4. For $g \geq 1$, it is easy to check that $\overline{a_{(g,(2g+1)p+g)}} = \overline{a_{(0,p)}}$. The Y-coordinate of $\overline{a_{(g,p+1)}}$ is greater than that of $\overline{a_{(g,p)}}$, because $\frac{2(p+1)+1}{2(p+1)+2g+2} > \frac{2p+1}{2p+2g+2}$. These imply that $\overline{a_{(g,(2g+1)p+i)}}$ for each $g \leq i \leq 3g+1$ lies between $\overline{a_{(0,p)}}$ and $\overline{a_{(0,p+1)}}$.

In particular $\lim\limits_{p \to \infty} \frac{p \log r_{(g,p)}}{\log p} = 1$.

Lemmas 4.4 and 4.5 below allow us to construct a $\Sigma_{g,2p+i}$-bundle over the circle for each $i \in \{1,2,3,4\}$ with the dilatation $r_{(g,p)}$, provided $a_{(g,p)}$ is primitive.

Lemma 4.4. *Suppose that $a_{(g,p)}$ is primitive. Then the minimal representative $F_{a_{(g,p)}}$ is a surface of genus g with $2p+4$ boundary components, and the stable foliation $\mathcal{F}_{a_{(g,p)}}$ has the following singularities.*

1. *If $p+g$ is odd, then $\sharp(\partial_\alpha F_{a_{(g,p)}}) = 2$, $\sharp(\partial_\beta F_{a_{(g,p)}}) = 2p+1$ and $\sharp(\partial_\gamma F_{a_{(g,p)}}) = 1$. In this case $\mathcal{F}_{a_{(g,p)}}$ is $\frac{p+g+1}{2}$-pronged at each component of $\partial_\alpha F_{a_{(g,p)}}$, 1-pronged at each component of $\partial_\beta F_{a_{(g,p)}}$ and $(p+3g+2)$-pronged at the component of $\partial_\gamma F_{a_{(g,p)}}$.*
2. *If $p+g$ is even, then $\sharp(\partial_\alpha F_{a_{(g,p)}}) = 1$, $\sharp(\partial_\beta F_{a_{(g,p)}}) = 2p+1$ and $\sharp(\partial_\gamma F_{a_{(g,p)}}) = 2$. In this case $\mathcal{F}_{a_{(g,p)}}$ is $(p+g+1)$-pronged at the component of $\partial_\alpha F_{a_{(g,p)}}$, 1-pronged at each component of $\partial_\beta F_{a_{(g,p)}}$ and $\frac{p+3g+2}{2}$-pronged at each component of $\partial_\gamma F_{a_{(g,p)}}$.*

In both cases (1) and (2), $\mathcal{F}_{a_{(g,p)}}$ has no singularities in the interior of $F_{a_{(g,p)}}$.

Proof. By definition, $a_{(g,p)} = (p+g+1)\mathfrak{a} + (p-q)\mathfrak{b}$. Notice that the parity of $p+g+1$ is different from that of $p-g$. By Lemma 4.1, $\sharp(\partial_\alpha F_{a_{(g,p)}}) + \sharp(\partial_\gamma F_{a_{(g,p)}})$ is equal to 3.

The singularity data of $\mathcal{F}_{a_{(g,p)}}$ is obtained from the formula in Section 4.2. □

By Lemma 4.4, it is straightforward to prove

Lemma 4.5. *Suppose that $a_{(g,p)}$ is primitive. Then $\mathcal{F}_{a_{(g,p)}}$ is no 1-pronged at each component of $\partial_\alpha F_{a_{(g,p)}} \cup \partial_\gamma F_{a_{(g,p)}}$ if and only if $(g,p) \notin \{(0,0),(0,1),(1,0)\}$. In particular if $g \geq 2$ and $p \geq 0$, then $\mathcal{F}_{a_{(g,p)}}$ is no 1-pronged at each component of $\partial_\alpha F_{a_{(g,p)}} \cup \partial_\gamma F_{a_{(g,p)}}$.*

We are now ready to prove theorems in Section 1.

Proof of Theorem 1.4. There exists a sequence of primitive fibered classes $\{a_{(g,p_i)}\}_{i=0}^{\infty}$ with $p_i \to \infty$. (In fact, if $p_i = (g+1)+(2g+1)i$, then $2g+1$ and p_i+g+1 are relatively prime, and hence $a_{(g,p_i)}$

is primitive.) Then N is a $\Sigma_{g,2p_i+4}$-bundle over the circle with the dilatation $r_{(g,p_i)}$. Therefore $\delta_{g,2p_i+4} \le r_{(g,p_i)}$. If we set $n_i = 2p_i + 4$, then

$$\frac{n_i \log \delta_{g,n_i}}{\log n_i} \le \frac{n_i \log r_{(g,p_i)}}{\log n_i} = \frac{(2p_i+4) r_{(g,p_i)}}{\log(2p_i+4)}.$$

By Lemmas 2.3 and 4.3, $\frac{(2p_i+4)r_{(g,p_i)}}{\log(2p_i+4)}$ goes to 2 as i goes to ∞. Therefore

$$\limsup_{i\to\infty} \frac{n_i \log \delta_{g,n_i}}{\log n_i} \le 2.$$

$\square$

Proof of Theorem 1.5. The minimal representative of $a_{(g,p)}$ is a surface of genus g with $2p+4$ boundary components with

$$\mathrm{slope}(a_{(g,p)}) = (\tfrac{3p-g+1}{-(p+q+1)}, -1, \tfrac{3p+g+2}{-(p-g)}).$$

Thus N is a $\Sigma_{g,2p+4}$-bundle over the circle with the dilatation $r_{(g,p)} = \lambda(a_{(g,p)})$, and hence $\delta_{g,2g+4} \le r_{(g,p)}$.

Consider the case where $p+g$ is odd. (See Lemma 4.4(1).) The manifold $N(\frac{3p+g+2}{-(p-g)})$, obtained from N by Dehn filling a cusp specified by the torus T_γ along the boundary slope $\frac{3p+g+2}{-(p-g)}$ of the fiber $F_{a_{(g,p)}}$, is a $\Sigma_{g,2p+3}$-bundle over the circle with the dilatation $r_{(g,p)}$ since $\mathcal{F}_{a_{(g,p)}}$ is no 1-pronged at each component of $\partial_\gamma F_{a_{(g,p)}}$. Similarly, by Lemma 4.5, we see that $N(\frac{3p-g+1}{-(p+q+1)})$ (resp. $N(\frac{3p-g+1}{-(p+q+1)}, \frac{3p+g+2}{-(p-g)})$) is a $\Sigma_{g,2p+2}$-bundle (resp. $\Sigma_{g,2p+1}$-bundle) over the circle with the dilatation $r_{(g,p)}$. The existence of these $\Sigma_{g,2p+i}$-bundles for $i \in \{1,2,3\}$ imply that $\delta_{g,2p+i} \le r_{(g,p)}$.

The case where $p+g$ is even is analogous. $\square$

Proof of Theorem 1.6. By Theorem 1.5 we have the following:

- If $\gcd(2g+1, p+g+1) = 1$ and $\gcd(2g+1, (p+1)+g+1) = 1$, then for $j \in \{3,4\}$
$$\delta_{g,2p+j} \le \min\{r_{(g,p)}, r_{(g,p+1)}\}.$$
- If $\gcd(2g+1, p+g+1) = 1$ and $\gcd(2g+1, (p+1)+g+1) \ne 1$, then for $j \in \{3,4\}$
$$\delta_{g,2p+j} \le r_{(g,p)}.$$
- If $\gcd(2g+1, p+g+1) \ne 1$ and $\gcd(2g+1, (p+1)+g+1) = 1$, then for $j \in \{3,4\}$
$$\delta_{g,2p+j} \le r_{(g,p+1)}.$$

Thus

$$\frac{(2p+j)\log \delta_{g,2p+j}}{\log(2p+j)} \le \frac{(2p+j)\log r_{(g,p)}}{\log(2p+j)} \quad \text{or} \quad \frac{(2p+j)\log \delta_{g,2p+j}}{\log(2p+j)} \le \frac{(2p+j)\log r_{(g,p+1)}}{\log(2p+j)}. \tag{4.1}$$

$(*)$ in Theorem 1.6 ensures that (4.1) holds for any $p \ge 0$. By using Lemma 2.3, it is easy to see that both $\frac{(2p+j)\log r_{(g,p)}}{\log(2p+j)}$ and $\frac{(2p+j)\log r_{(g,p+1)}}{\log(2p+j)}$ go to 2 as p goes to ∞. Thus

$$\limsup_{p\to\infty} \frac{(2p+j)\log \delta_{g,2p+j}}{\log(2p+j)} \le 2,$$

which means that

$$\limsup_{n\to\infty} \frac{n \log \delta_{g,n}}{\log n} \le 2.$$

$\square$

Proof of Proposition 1.10. We prove the claim in the second half. (The proof in the first half is similar.) If $g \geq 2$ satisfies $(*)$, then for any $p \geq 0$ there exist a $\Sigma_{g,2p+3}$-bundle and a $\Sigma_{g,2p+4}$-bundle over the circle obtained from N, see Proof of Theorem 1.6. More precisely such a bundle is homeomorphic to N or it is obtained from N by Dehn filling cusps along the boundary slopes of a fiber. Thus Proposition 1.10 holds from the result which says that hyperbolic volume decreases after Dehn filling [16, 18]. □

Appendix A. Examples by Tsai

Given $g \geq 2$, Tsai gave in [22] the examples of pseudo-Anosovs $f_{g,n}$ defined on an orientable surface of genus g with n marked points which enjoy $\log \lambda(f_{g,n}) \asymp \frac{\log n}{n}$. By puncturing the marked points, one can obtain the pseudo-Anosov mapping class on $\Sigma_{g,n}$, a surface of genus g with n punctures with the same dilatation as $f_{g,n}$. Following [22], we briefly recall her construction of $f_{g,n}$ on a surface of genus g with n marked points, and we denote such a surface by $\Sigma_{g,n}$ by abusing the notation again. Then we prove

Proposition A.1. *For any fixed* $g \geq 2$, $\displaystyle\lim_{n\to\infty} \frac{n \log \lambda(f_{g,n})}{\log n} = 2(2g+1)$.

For $p, q \geq 1$, we now define a mapping class $\phi_{p,q} \in \mathrm{Mod}(\Sigma_{0,p+q+2})$. First, let $\Sigma_{0,p+q+2}$ be a sphere with $(p+1)$ marked points X circling unmarked point x, and $(q+1)$ marked points Y circling unmarked point y, and a single marked point z, see Figure 5(left). Note that $|X \cap Y| = 1$, $|X| = p+1$ and $|Y| = q+1$. Second, define homeomorphisms $\alpha_p, \beta_q : \Sigma_{0,p+q+2} \to \Sigma_{0,p+q+2}$ so that α_p rotates the marked points of X counterclockwise around x and β_q rotates the marked points of Y clockwise around y, see Figure 5(center, right). Then let $\phi_{p,q} = [\beta_q] \circ [\alpha_p]$. This mapping class $\phi_{p,q}$ is pseudo-Anosov, and $\lambda(\phi_{p,q})$ is the largest real root of

$$T_{p,q}(t) = t^{p+q+1}(t-1) + 1 - 2(t^{p+1} + t^{q+1}) - t,$$

see [7]. By Theorem 2.1, $\lambda(\phi_{p,q})$ is indeed a unique real root greater than 1 of $T_{p,q}(t)$. The following property holds, see [10, Proposition 3.4]: For $p' > p$ and $q' > q$,

$$\lambda(\phi_{p',q}) < \lambda(\phi_{p,q}) \text{ and } \lambda(\phi_{p,q'}) < \lambda(\phi_{p,q}). \tag{A.1}$$

The lemma (see [10, Lemma 2.2]) on the theory of Perron-Frobenius matrices was used for the proof of these inequalities.

It was shown in [7] that

$$\lambda(\phi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}) = \min\{\lambda(\phi_{p,q}) \mid p+q = s\}.$$

Tsai proved that $\log \lambda(\phi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}) \asymp \frac{\log s}{s}$, see [22, Proposition 4.1]. More generally if we fix $k, \ell \geq 0$, then $\log \lambda(\phi_{p+k,p+\ell}) \asymp \frac{\log p}{p}$ holds. In fact we have the following lemma which is a key to prove Proposition A.1.

Lemma A.2. *We fix* $k, \ell \geq 0$. *Given* $0 < c_1 < 1$ *and* $c_2 > 1$, *there exists* $L > 0$ *such that*

$$p^{\frac{c_1}{p}} < \lambda(\phi_{p+k,p+\ell}) < p^{\frac{c_2}{p}} \quad \textit{for any } p > L.$$

Proof. The dilatation $\lambda(\phi_{p+k,p+\ell})$ is a unique real root greater than 1 of

$$T_{p+k,p+\ell}(t) = t^{2p+k+\ell+1}(t-1) + 1 - 2(t^{\ell+1} + t^{k+1})t^p - t.$$

The lemma follows from Proposition 2.2(2). □



FIGURE 5. (left) sphere with marked points $\Sigma_{0,p+q+2}$ ($p=q=5$), (center, right) homeomorphisms α_p and β_q.



FIGURE 6. invariant train track for $\phi_{p,q}$ ($p=q=5$).



FIGURE 7. images of the train track under $\alpha_{p,q}$ (left) and under $\beta_{p,q}$ (right), ($p=q=5$).

We recall the pseudo-Anosov $f_{2,n} \in \mathrm{Mod}(\Sigma_{2,\mathrm{n}})$, and then we prove Proposition A.1 in the case $g=2$. Let $s=p+q$.

We can construct the cyclic branched cover $\Sigma_{2,n}$ of $\Sigma_{0,s+2}$ with branched points x, y, z, where $n = 5(s+1)+1$ so that both $\alpha_p, \beta_q : \Sigma_{0,s+2} \to \Sigma_{0,s+2}$ are lifted to $\widetilde{\alpha}_p, \widetilde{\beta}_q : \Sigma_{2,n} \to \Sigma_{2,n}$, see [22, Section 4.1]. One can check that both $\alpha_p, \beta_q : \Sigma_{0,s+2} \to \Sigma_{0,s+2}$ can be lifted to $\widetilde{\alpha}_p, \widetilde{\beta}_q : \Sigma_{2,n} \to \Sigma_{2,n}$. Define $\psi_{p,q} = [\widetilde{\beta}_q] \circ [\widetilde{\alpha}_p] \in \mathrm{Mod}(\Sigma_{2,n})$. The mapping class $\psi_{p,q}$ is pseudo-Anosov with $\lambda(\psi_{p,q}) = \lambda(\phi_{p,q})$, since $\psi_{p,q}$ is a lift of $\phi_{p,q}$.

By taking $p = \lfloor \frac{s}{2} \rfloor$ and $q = \lceil \frac{s}{2} \rceil$, we obtain $\psi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil} \in \mathrm{Mod}(\Sigma_{2,5(s+1)+1})$. Tsai modified $\psi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}$ to be a pseudo-Anosov mapping class $\psi_{(s;i)} \in \mathrm{Mod}(\Sigma_{2,5(s+1)+1+i})$ for each $i \in \{0,1,2,3,4\}$ so that $\psi_{(s;0)} = \psi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}$. Moreover she proved that

$$\lambda(\psi_{(s;i)}) \le \lambda(\psi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}) \tag{A.2}$$

by using [22, Proposition 2.5] on the theory of the Perron Frobenius matrices. One can verify that the inequality

$$\lambda(\psi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) \leq \lambda(\psi_{(s;i)})$$

holds by using the same argument as in the proof of (A.1).

Putting the two inequalities above together, we have

$$\lambda(\psi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) \leq \lambda(\psi_{(s;i)}) \leq \lambda(\psi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}).$$

Let $m = \lfloor \frac{s}{2} \rfloor$. Given $0 < c_1 < 1$ and $c_2 > 1$, there exists $L_i > 0$ for each $i \in \{0, 1, 2, 3, 4\}$ such that

$$m^{\frac{c_1}{m}} < \lambda(\phi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) < m^{\frac{c_2}{m}} \quad \text{for } m > L_i,$$

see Lemma A.2. Thus

$$m^{\frac{c_1}{m}} < \lambda(\phi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) < m^{\frac{c_2}{m}} \quad \text{for } m > L = \max\{L_i \mid i \in \{0, 1, 2, 3, 4\}\}.$$

Since $\lambda(\psi_{p,q}) = \lambda(\phi_{p,q})$,

$$m^{\frac{c_1}{m}} < \lambda(\phi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) \leq \lambda(\psi_{(s;i)}) \leq \lambda(\phi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}) < m^{\frac{c_2}{m}} \quad \text{for } m > L.$$

For each n, let $f_{2,n} = \psi_{(s;i)}$, where s and $i \in \{0, 1, 2, 3, 4\}$ satisfy $n = 5(s+1)+1+i$. By Lemma 2.3 together with $m = \lfloor \frac{s}{2} \rfloor$, we have

$$\lim_{n \to \infty} \frac{n \log \lambda(f_{2,n})}{\log n} = \lim_{s \to \infty} \frac{(5(s+1)+1+i) \log \lambda(\psi_{(s;i)})}{\log(5(s+1)+1+i)} = 10.$$

This completes the proof in the case $g = 2$.

Tsai generalized her construction of $f_{2,n} \in \mathrm{Mod}(\Sigma_{2,n})$ to $f_{g,n} \in \mathrm{Mod}(\Sigma_{g,n})$ for any fixed $g > 2$ as we briefly recall below. The pseudo-Anosov mapping class $\psi_{(g,s)} \in \mathrm{Mod}(\Sigma_{g,n})$ can be defined by taking a suitable branched cover over $\Sigma_{0,s+2}$, where $n = (2g+1)(s+1)+1$. Then she extended to arbitrary n by constructing $\psi_{(g,s;i)} \in \mathrm{Mod}(\Sigma_{g,(2g+1)(s+1)+1+i})$ for each $i \in \{0, 1, \cdots, 2g\}$ so that $\psi_{(g,s)} = \psi_{(g,s;0)}$. By using the same argument as above, one obtains

$$\lambda(\phi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) \leq \lambda(\psi_{(g,s;i)}) \leq \lambda(\phi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}).$$

Furthermore given $0 < c_1 < 1$ and $c_2 > 1$, there exists $L > 0$ such that

$$m^{\frac{c_1}{m}} < \lambda(\phi_{\lfloor \frac{s}{2} \rfloor + i, \lceil \frac{s}{2} \rceil}) \leq \lambda(\phi_{\lfloor \frac{s}{2} \rfloor, \lceil \frac{s}{2} \rceil}) < m^{\frac{c_2}{m}} \text{ for } m > L,$$

where $m = \lfloor \frac{s}{2} \rfloor$. Therefore

$$m^{\frac{c_1}{m}} < \lambda(\psi_{(g,s;i)}) < m^{\frac{c_2}{m}} \text{ for } m > L.$$

For each n, let $f_{g,n} = \psi_{(g,s;i)}$, where s and $i \in \{0, 1, \cdots, 2g\}$ satisfy $n = (2g+1)(s+1)+1+i$. By Lemma 2.3 we have

$$\lim_{n \to \infty} \frac{n \log \lambda(f_{g,n})}{\log n} = 2(2g+1).$$

Thus we have obtained Proposition A.1.

References

[1] J. W. Aaber and N. M. Dunfield, *Closed surface bundles of least volume*, Algebraic and Geometric Topology 10 (2010), 2315-2342.

[2] A. Casson and S. Bleiler, Automorphisms of surfaces after Nielsen and Thurston, Cambridge University Press (1988).

[3] J. H. Cho and J. Y. Ham, *The minimal dilatation of a genus-two surface*, Experimental Mathematics 17 (2008), 257-267.

[4] D. Fried, *Flow equivalence, hyperbolic systems and a new zeta function for flows*, Commentarii Mathematici Helvetici 57 (1982), 237-259.

[5] J. Y. Ham and W. T. Song, *The minimum dilatation of pseudo-Anosov* 5*-braids*, Experimental Mathematics 16 (2007), 167-179.

[6] E. Hironaka, *Small dilatation mapping classes coming from the simplest hyperbolic braid*, Algebraic and Geometric Topology 10 (2010), 2041-2060.

[7] E. Hironaka and E. Kin, *A family of pseudo-Anosov braids with small dilatation*, Algebraic and Geometric Topology 6 (2006), 699-738.

[8] N. V. Ivanov, *Stretching factors of pseudo-Anosov homeomorphisms*, Journal of Soviet Mathematics, 52 (1990), 2819–2822, which is translated from Zap. Nauchu. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 167 (1988), 111–116.

[9] E. Kin, S. Kojima and M. Takasawa, *Minimal dilatations of pseudo-Anosovs generated by the magic* 3*-manifold and their asymptotic behavior*, preprint (2011), arXiv:1104.3939v2

[10] E. Kin and M. Takasawa, *An asymptotic behavior of the dilatation for a family of pseudo–Anosov braids*, Kodai Mathematical Journal 31 (2008), 92-112.

[11] E. Kin and M. Takasawa, *Pseudo-Anosov braids with small entropy and the magic* 3*-manifold*, Communications in Analysis and Geometry 19 (4) (2011), 705-758.

[12] E. Kin and M. Takasawa, *Pseudo-Anosovs on closed surfaces having small entropy and the Whitehead sister link exterior*, preprint (2010), arXiv:1003.0545, to appear in "Jounal of the Mathematical Society of Japan".

[13] K. H. Ko, J. Los and W. T. Song, *Entropies of braids*, Journal of Knot Theory and its Ramifications 11 (2002), 647-666.

[14] E. Lanneau and J. L. Thiffeault, *On the minimum dilatation of braids on the punctured disc*, Geometriae Dedicata 152 (2011), 165-182.

[15] C. McMullen, *Polynomial invariants for fibered* 3*-manifolds and Teichmüler geodesic for foliations*, Annales Scientifiques de l'École Normale Supérieure. Quatrième Série 33 (2000), 519-560.

[16] W. D. Neumann and D. Zagier, Volumes of hyperbolic three-manifolds, Topology 24 (3) (1985), 307-332.

[17] R. C. Penner, *Bounds on least dilatations*, Proceedings of the American Mathematical Society 113 (1991), 443-450.

[18] W. Thurston, The geometry and topology of 3-manifolds, Lecture Notes, Princeton University (1979).

[19] W. Thurston, *A norm of the homology of* 3*-manifolds*, Memoirs of the American Mathematical Society 339 (1986), 99-130.

[20] W. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bulletin of the American Mathematical Society 19 (1988), 417-431.

[21] W. Thurston, *Hyperbolic structures on 3-manifolds II: Surface groups and 3-manifolds which fiber over the circle*, preprint, arXiv:math/9801045

[22] C. Y. Tsai, *The asymptotic behavior of least pseudo-Anosov dilatations*, Geometry and Topology 13 (2009), 2253-2278.

[23] C. Y. Tsai, *Minimal pseudo-Anosov translation lengths on the Teichmüler space*, PhD thesis, University of Illinois at Urbana-Champaign (2010).

[24] A. D. Valdivia, *Sequences of pseudo-Anosov mapping classes and their asymptotic behavior*, preprint (2012), arXiv:1006.4409v3

[25] X. Wang, *A simple proof of Descartes's rule of signs*, American Mathematical Monthly 111 (6) (2004), 525-526.

Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Ohokayama, Meguro, Tokyo 152-8552 Japan
E-mail address: `kin@is.titech.ac.jp`

Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Ohokayama, Meguro, Tokyo 152-8552 Japan
E-mail address: `takasawa@is.titech.ac.jp`